2017

www.jmscr.igmpublication.org Impact Factor 5.84 Index Copernicus Value: 83.27 ISSN (e)-2347-176x ISSN (p) 2455-0450 crossref DOI: https://dx.doi.org/10.18535/jmscr/v5i2.12



Journal Of Medical Science And Clinical Research An Official Publication Of IGM Publication

Varied Modalities in Management of Fractures in Patients with Post Polio Residual Paralysis-Case Series

Authors

Dr T.Sathish Kumar¹, Prof. Dr A.Senthilnathan², Dr R.Prabhakar³

¹Postgraduate in Orthopaedic Surgery, RMMCH, Annamalai University ²Professor & Chief, Department of Orthopaedic Surgery, RMMCH, Annamalai University ³Lecturer, Department of Orthopaedic Surgery, RMMCH, Annamalai University

Abstract

Introduction: Poliomyelitis is on the verge of eradication. Fractures in polio survivors present unique challenges; the bone is often small, deformed, osteoporotic and hypovascularized.

Aim: Our aim was to assess the varied fracture patterns and the challenges faced in fixation of fractures in polio affected limbs and possible solutions to overcome them.

Material & Methods: All cases were managed at RMMCH, Annamalai University. Each and every case was unique in its own terms with variable clinical and radiological presentation. Varied challenges were managed in different manner with respect to implant choice for optimal outcome.

Results: Fracture union and callus formation were evaluated on radiographs taken at each postoperative visit. Functional outcome assessments included range of motion and early return to activities of daily living. All fractures healed except for one non union due to conservative management.

Conclusion: Anatomical locking plate may not be ideal for these patients as the plates may be larger and overhang the malformed bone. Hence nails with smaller diameter and locking compression plates designed for other skeletal regions may give better results than the conventional plates and nails. Fractures in the polio-affected limb are not a common entity and are difficult to manage using a common fracture management protocols and need to be customized for individual patients to suit their needs and bones with all implants ready for use.

Keywords: Post-polio residual paralysis, Fracture Fixation.

INTRODUCTION

Poliomyelitis is on the verge of eradication. But the survivors of polio are still living with its consequences in different parts of the world. People with post-polio residual paralysis are prone to fractures after mild trauma. The flaccid paralysis, asymmetric involvement, and underdeveloped growth of affected leg may lead to osteoporosis which is the common cause for fracture. Fractures in polio survivors present unique challenges; the bone is often small, deformed, osteoporotic and hypovascularized.

AIM

Our aim is to assess the varied fracture patterns and the challenges faced in fixation of fractures in polio affected limbs and possible solutions to overcome them. Knowledge of treating these fractures is important to a trauma surgeon as such scenarios are not common in daily practice.

MATERIAL & METHODS

All 7 cases were managed by the Department of Orthopaedic Surgery, Rajah Muthiah Medical College & Hospital, Annamalai University, Chidambaram. Each and every case was unique in its own terms with variable clinical and radiological presentation and varied challenges & were managed in a different manner with respect to implant choice for optimal outcome.

DISCUSSION

Case 1: 30 year old male presented with h/o of self fall (low energy trauma) with PPRP





Pre Operative Xray Showing Fracture Of Proximal Shaft Right Femur With Intercondylar Fracture

Options available for Fixation

- Supra condylar nail with intercondylar cc screws
- narrow dcp
- distal femur locking plate.
- 95 degree angle blade plate for intercondylar fracture.

Our Plan & Treatment

Patient was not using the fractured limb for weight bearing & mobilization prior to injury. Hence our original plan was to fix with a distal femur locking compression plate. Per operatively we found that even the smallest plate was much larger than the deformed bone. Instead we Used a Philos plate for fixation of intercondylar fracture in distal femur (part of segmental fracture) and dynamic compression plate (narrow dcp) for ipsilateral shaft of femur fracture in patient with fixed flexion & rotational deformity of the affected knee



Post –op X rays showing philos plate for distal femur with intercondylar extension & narrow dcp for proximal shaft of femur right

Advantages of using philos plate

- It is a relatively strong implant.
- Multiple screws in varied direction gives very strong fixation.
- Can be considered as an alternative where distal femur locking compression plate are larger than the bone and multiple purchase are required in the fracture fragment

Case 2: 59 year old male presented with h/o of self fall (low energy trauma) with PPRP



Pre Operative Xray Showing Supra Condylar Fracture Distal Femur Right Side

Options available for Fixation

- Supra condylar nail
- Distal femur locking plate⁷.
- DYNAMIC CONDYLAR PLATE
- PHILOS plate

Our Plan & Treatment

Patient was active and mobile using the fractured limb for weight bearing & mobilization prior to injury (hand on knee gait). Our plan was to use either a distal femur locking compression plate or a PHILOS plate as we did in our earlier case. But on the table we used a Proximal tibia locking compression plate⁵ (anterolateral) for supra condylar fracture distal femur which seated well on the bone without over hanging of the implant & also gave adequate fixation.



Post –op X rays showing proximal tibia LCP for supracondylar femur fracture

Advantages of using proximal tibia LCP plate

- Relatively smaller implant compared to distal femur lcp.
- Adequate screws in distal fragment gives very strong fixation.

Case 3: 30 year old male presented with h/o of RTA with bilateral



Pprp Pre Operative Xray Showing Supra Condylar Fracture Distal Femur Right Side

Options available for Fixation

- Supra condylar nail
- Distal femur locking plate⁷
- PHILOS plate

Our Plan & Treatment

Patient was not using the fractured limb for weight bearing & mobilization prior to injury. Patient was using his upper limb for active mobilization. Our plan was to use a proximal tibia lcp which we used for our previous case but the Proximal tibia locking compression plate⁵ (POSTEROMEDIAL) was more apt than the anterolateral plate for supra condylar fracture distal femur which gave good fixation without any hardware problem⁷.



Post –op X rays showing proximal tibia LCP for supracondylar femur fracture

Advantages of using proximal tibia LCP

- Implant provides more distal fixation compared to anterolateral lcp.
- Shape of the plate was more apt for the deformed bone than the anterolateral plate.

Case 4: 40 year old male presented with h/o of RTA and injury to left thigh with PPRP



Pre Operative Xray Showing fracture shaft of femur left side with another fracture line extending from trochanteric - subtrochanteric region

Options available for Fixation

- Broad dcp
- 95 degree angle blade plate with long side plate^{1,2,3}
- Long PFN

Our Plan & Treatment

Patient was active and mobile using the fractured limb for weight bearing & mobilization prior to injury (hand on knee gait).Our plan was to use a 95 degree angle blade plate with long side plate using MIPO technique .But we Used an Intramedullar Interlocking nail (Closed reduction) for fracture shaft of femur with a Locking bolt(extra medullary) as lag screw to address the trochanteric- subtrochanteric (reverse Oblique) extension.



Post –Op X Rays Showing Imil Nail With Lag Screw In Sub Trochanteric Region

Advantages of using Intramedullar Interlocking nail

• Implant of choice for fixation of long bones.

2017

- Early mobilization & weight bearing
- Provides stability in all planes

Case 5: 49 year old male presented with h/o of RTA and injury to Right thigh with PPRP - presented to us three weeks after injury. Had initially taken native treatment in the form of splints which were changed weekly for 3 weeks



Pre Operative Xray Showing Fracture Shaft Of Femur Right Side

Options available for Fixation

- Broad/Narrow DCP dcp
- K nail
- Small size (7,8 size)IMIL nail

Our Plan & Treatment

Patient was active and mobile using the fractured limb for weight bearing & mobilization prior to injury. Our plan was to use a IMIL nail of smaller diameter. However we also had regular size nails taking into account the wide medullary canal. We used a 9x40 size Intramedullar Interlocking nail (closed reduction) with 2 distal locking bolts and 1 proximal locking bolt as we were able to use the 10 size reamer with ease.



Post –Op X Rays Showing Imil Nail With Locking Bolts

2017

Advantages of using Intramedullar Interlocking nail

- Implant of choice for fixation of long bones.
- Early mobilization & weight bearing
- Provides stability in all planes

Case 6: 65 year old female sustained self fall (low velocity injury) and injured right hip



Pre Operative Xray Showing Fracture Neck of Femur Right Side

Options available for Fixation

- ORIF with cc screws
- ORIF with DHS
- Primary THA^6

Our Plan & Treatment

Patient was house wife and requested for early ambulation to perform regular house hold activities. We anticipated difficulties in management of fracture neck of femur in a paralytic limb due to excessive anteversion, valgus neck, small neck, dislocated and dysplastic hip⁴. We planned for hemiarthroplasty considering it as one step solution for early weight bearing taking into account the possibility of failure and the need for second surgery. We have ordered bipolar prosthesis with long stem anticipating various difficulties but we used the Austin moore prosthesis with regular stem and started weight bearing on 3rd post operative day and patient walked back home on the 13th Post Operative day.



Post –Op X Rays Showing Uncemented Hemiarthroplasty Using Austin Moore Prosthesis

Advantages of doing Uncemented Hemiarthroplasty

- 1. Cost effective
- 2. Early Weight bearing rather than CRIF/ORIF with CC screws/DHS
- 3. Can plan for revision THA at a late date if required

Case 7: 43 year old male sustained injury due to RTA and injured left leg 6 months ago-took native treatment for 3 months and presented to us with the below mentioned x ray.



Initial Xray Showing (3 Months Old) Spiral Fracture of Lower Third Of Tibia Left

Options available for Fixation

- Distal tibia lcp
- Recon/1/3 tubular plate for fibula

Our Plan & Treatment

We planned for open reduction and internal fixation with distal tibia locking compression plate and 1/3 tubular plate for fibula .Patient has developed aversion for surgery as he had undergone multiple surgeries during childhood. Patient was already walking comfortably with calipers and came to the OPD just to know if fracture has united. Hence conservative manage-

2017

ment for this patient who refused to undergo surgery in distal tibia fracture



Recent X Ray Showing (6 Months Old) Spiral Fracture Of Lower Third Of Tibia Left (Delayed Union)

Disadvantages of Conservative therapy

- 1. Delayed Union going in for non union
- 2. Osteoporosis due prolonged immoblisation
- 3. No adequate callus formation

RESULTS

All fractures maintained fairly good alignment post operatively. Fracture union and callus formation were evaluated on radiographs taken at each postoperative visit at 3,6,12,18 weeks. Functional outcome assessments included range of motion and early return to activities of daily living. All fractures healed except for one non union due to conservative management. But union was delayed. There were no cases of implant cutout, failure or other complications.

CONCLUSION

Early fixation and mobilization is optimal for patients with post polio residual paralysis as prolonged immobilization with cast would worsen osteopenia in already weak and deformed bones. Anatomical locking plate may not be ideal for these patients as the plates may be larger and overhang the malformed smaller osteoporotic bone. Conventional nails might be larger and broader resulting in iatrogenic fractures. Hence nails with smaller diameter and locking compression plates designed for other skeletal regions may give better results than the conventional plates and nails. Fractures in the polio-affected limb are not a common entity and are difficult to manage using a common fracture management protocols and need to be customized for individual patients to suit their needs and bones with all armaments (implants) ready for use.

REFERENCES

- Journal of Evolution of Med and Dent Sci/ eISSN- 2278-4802, pISSN- 2278-4748/ Vol. 3/ Issue 61/Nov 13, 2014
- Muller ME, Allgower M, Schneider R et al. Manual of Internal Fixation, 3rd Ed. Berlin: Springer Verlag, 1990.
- 3. Kinast C, Bolhofner BR, Mast JW and Ganz R. Subtrochanteric fractures of the femur: results of treatment with the 95" condylar blade plate. Clin Orthop 1989; 238: 122.
- Journal of Evolution of Medical and Dental Sciences 2014; Vol. 3, Issue 30, July 28;
- Pub Med: Archives of Orthopaedic and Trauma Surgery 130(10):1299-304October 2010
- Hip international: the journal of clinical and experimental research on hip pathology and therapy 21(2):267-269 · April 2011.
- 7. Distal femoral fractures in postpoliomyelitis patients treated with locking compression plates. Wang WJ, Shi HF, Chen DY, Chen YX, Wang JF, Wang SF, Qiu Y, Xiong J. 2013 Chinese Association Orthopaedic and Wiley Publishing Asia Pty Ltd.